

Fungi Lusitani

- 1 *Phoma herbarum* West. forma
- 2 *Phoma Allioniae* Bres. n. sp.
 Peritheciis dense gregariis, primo subepidermicis, dein cortice sec-
 dente demudatis, globoso-subdepressis, ostiolo proclivis 100-140 μ .
 diam., membranaceis, contentu prosenchymatico, luride castoreo;
 sporulis elongatis, hyalinis, 4-6 \times 2-2 1/2 μ . —
 Hab. in caulibus *Allioniae violaceae* L., Coimbra
 Herb. Botanicis, Nov. 1897 (C. Heller).
- 3 *Ximonium velutatum*, indeterminabile
4. *Phyllosticta Glycines* Thüm. et *Gleosporium intermedium* Sacc.
 (species ultime in foliis supra albicis)
5. *Coccomyces trigony* Thüm. non K. et Schm. qui est forma
Coccomyces turgide Fr.
6. *Gyrocera* Celtidis (Bivona) M. et Ces.
7. Nihil vivens — non fungus —
8. Imaturus — indeterminabile —
9. *Phyllachora Ulmi* (Dav.) Fückel
10. Imaturus — indeterminabilis —

11. Nihil inveni
12. Physalospora laligans Sacc. n. sp. ff. 12 et ff. 20
 in dubio ne cum speciebus a Saccardo in sua "Contributio
 Florae myc. lusitanicae", concordant, Saccardo misit, et ab
 eo prope cum novae speciei recognoscitae, scripsit, quod
 Sagnosus ipse statim Clar. Prof. Peniquez in calce
 operis sui misit. Hic ergo querendae —
13. Phoma viminalis Cooke
14. Nihil inveni —
15. Harknessia uromycoidea Speg.
16. Nimium vetustas, indeterminabilis
17. Id. " "
18. Marasmius hygrometricus Briganti.
19. Imaturus, indeterminabilis
20. Phoma Achilleae Sacc. * Ph. Dahliae Sacc. n. subsp.
21. Coccomyces dentatus (Kunze et Schm.) Sacc.
22. Phoma Rusci West —
23. Phoma Oleae (DeC.) Sacc.

24. Imature, indeterminabiliz
25. Nihil inveni nisi maculas
26. Non fungus —
27. Imature, forsan Phyllosticta Arbuti-umedonic
28. Nihil inveni —
29. Cercospora tinca Sacc. —
30. Cercospora Smilacis Thüm. —

